

MAIN FILE

OTS: 60-41,660

JPRS: 4115

MAIN FILE

27 October 1960

THE WORK OF THE DIVISION OF IMMUNOLOGY AND ONCOLOGY  
OF THE INSTITUTE OF EPIDEMIOLOGY AND MICROBIOLOGY

IMENI N. F. GAMALEYA

By L. A. Zil'ber

- USSR -

TRANSFERRED TO MAIN FILE

19981208 057

Reproduced From  
Best Available Copy

DTIC QUALITY INSPECTED 2

Distributed by:

OFFICE OF TECHNICAL SERVICES  
U. S. DEPARTMENT OF COMMERCE  
WASHINGTON 25, D. C.

~~Price \$9.50~~

U. S. JOINT PUBLICATIONS RESEARCH SERVICE  
1636 CONNECTICUT AVE., N. W.  
WASHINGTON 25, D. C.

JPRS: 4115  
CSO: 1069-S/c

THE WORK OF THE DIVISION OF IMMUNOLOGY AND ONCOLOGY  
OF THE INSTITUTE OF EPIDEMIOLOGY AND MICROBIOLOGY  
IMENI N. F. GAMALEYA

-USSR-

[Following is the translation of an article entitled "O rabote Otdela immunologii i onkologii Instituta epidemiologii i mikrobiologii imeni N. F. Gamalei" (English version above) by L. A. Zil'ber in Vestnik Akademii Meditsinskikh Nauk (Herald of the Academy of Medical Sciences), Vol. 15, No. 7, Moscow, 1960, pages 65-72.]

The work of this Division is dedicated to the study of problems of general immunology, immunology of tumors and their etiology. This brief survey gives only the basic data obtained in the course of the last 2-3 years.

Among the problems of general immunology, a study of the question of a possible differentiation of the immune and the normal globulins was conducted (A. M. Gardash'yan and Z. A. Avenirova). For this purpose gamma-globulins were obtained from the serum of rabbits taken before and after immunization. Gamma globulins of the serum of the same rabbit were compared in a number of experiments. The purity of the preparations was checked electrophoretically. The comparison was done by the method of the cross anaphylaxis reaction with desensitization. These experiments showed an obvious difference in the antigenic structure of the normal and the immune globulin. Thus, the disturbance of the synthesis of the protein caused by the antigen in the process of immunization leads to the formation of globulins which differ in their antigenic structure from the globulins which are synthesized under normal conditions.

Antigenic structure of animal cells was studied in a number of works (G. I. Abelev, Z. A. Avenirova, Z. I. Baidakova, and N. V. Engel'garët). A study of this microsome, the mitochondria and the cell nuclei of the liver of mice has shown that the antigens of mitochondria and microsomes

are very close to each other, but are different from the antigens of the nucleus. The separation of organ-specific antigen of the liver of mice (AO) is of particular interest in the series of these studies. The principle of such separation is based on the fact that AO is bound more stably with an almost insoluble fraction of mitochondria and microsomes (Mm) than the rest of the antigens, and, therefore, can be separated from them.

When the suspension of Mm is destroyed with alkalizing or desoxycholate (0.3 percent), the main bulk of the Mm substance (and the greater part of antigens) changes to a solution, while a considerable part of AO remains bound with the undissolved fragments of Mm which settle in 60 minutes at 100 000 g. After the sediment was washed three times with a 0.3 percent desoxycholate, the antigen was changed into a solution by alkalizing.

A final purification was done with zonal electrophoresis in the density gradient, when AO, which possesses a very low electrophoretic mobility, remained near the point of the zone boundary. At the same time a fraction of non-antigenic lipoproteid which constituted more than 90 percent of the protein of the initial preparation, as well as antigenic admixtures, were separated from it.

The purified AO yielded one clear zone of precipitation in the agar with serums against liver and did not react with antihepatoma serums both in the precipitation reaction in the agar and in the reaction of quantitative precipitation. This antigen possessed a sharply expressed organ specificity in the precipitation reaction and was not evident among the antigens of the Mm of the kidneys, spleen, lungs, and plasma of mice.

The neutralization of the anti-liver serum with a fraction of Mm of the hepatoma and the inclusion of the antigens of the hepatoma into the agar did not hinder the reaction of AO with this serum. AO was found not only in the liver of mice of the C<sub>3</sub>HA strain, but also in the liver of mice C<sub>57</sub>, CC<sub>57</sub> and mice having no pedigree. At the present time, the chemical nature of this antigen is being studied.

Along with the antigenic structure of normal animal cells, the antigenic structure of tumor cells was studied. Investigations in the preceding years have established the

presence of antigens in tumor cells which are absent in normal cells (L. A. Zil'ber, 1950, 1957).

The success of these investigations was connected chiefly with the use of the anaphylaxis reaction with desensitization. Lately, the antigenic structure of tumor cells has been studied with the use of the precipitation reaction in the agar of the hepatoma model of C<sub>3</sub>HA mice induced by orthoazobenzidine and transplanted to mice of the same strain (G. I. Abelev, Z. A. Avenirova, Z. L. Baidakova, N. V. Engel'gardt).

Unlike the former works of our laboratory, cytoplasmic granules (Mm) were investigated, and not the protein fractions of tissue precipitated at pH 4.5, because our data and the data of other researchers point to antigenic changes of the granules during malignization. Proteins of other fractions were also studied. For a comparative evaluation of the results obtained by different methods, the anaphylaxis reaction with desensitization was also used. Used as antigens were solutions of Mm obtained by alkalizing their suspension to pH 9.0-9.5 with subsequent acidification to pH 7.8-8.2, as well as the extract of granules on 0.14 M NaCl.

Antiserums were prepared by immunizing rabbits against Mm and homogenates of normal liver and of hepatoma. Globulins of these serums were used in the precipitation reactions in the gel which was set up in the usual manner.

In order to expose more clearly the qualitative and quantitative differences of the preparations which were being compared, we applied the following arrangement of the reaction: Homologous serum and antigen are placed opposite one another in the opposite corners of a square. Antigens which are common for the systems which are being compared produce a single spectrum situated between the cavities with the heterologous antigen and the serums. Antigens which are characteristic for only one system produce bands directed along the diagonal of the square formed by the cavities with the antigens and the serums. These bands cd and ef are situated between the cavities with the antigens and the corresponding antiserum and their ends rest on the reservoirs of the heterologous system.

In the arrangement of the experiment described above which was suggested by G. I. Abelev, antigens characteristic

only for one of the systems form bands which differ in their direction from the general spectrum, and antigens and spectrum of two systems are compared here simultaneously. With such a method it was possible to expose qualitative differences between the antigenic structure of the tumor and the normal cells and to find an antigen which is specific for the hepatoma and which is absent in liver. This antigen has also been discovered during the neutralization of anti-hepatoma serums with Mm antigens of liver.

Recently, it became possible to separate a fraction of antibodies to the antigen of transplantable hepatoma which is absent in liver. Antibodies were separated by means of decomposing a precipitate obtained from antihepatoma serum neutralized by the excess of liver antigens during its reaction with the Mm of the hepatoma. The specific antibodies which were obtained facilitate the works on the separation of specific antigen of the hepatoma and the study of its nature.

These studies revealed a simplification of the antigenic structure of the cells during malignization which was described by Weiler (1952-1956) and Seligman, Grabar, Bernard (1955). The photograph shows precipitation lines which indicate that transplantable hepatoma contains an antigen which is absent in liver and at the same time does not have some of the liver antigens. In studying by the same method the nature of antigens which disappear during malignization, it was established that organ-specific antigen of liver (AO) is absent in the Mm of hepatoma, as it was also observed by Weiler in a rat hepatoma. However, anaphylaxis reaction with desensitization revealed an organ-specific antigen in hepatoma extracts containing, apart from Mm, proteins from other fractions (V. I. Gel'shtein). This problem is to be studied further.

Among the studies on the subject of immunology of tumors, we should note the data obtained by V. A. Parnes and D. M. Levina in collaboration with Dr. V. Lacour of the Gustave Roussie Institute in Paris, who worked with us. In agreement with the former data of our staff, V. V. Gordilova and L. V. Shersul'skaya, the antigenic difference of erythrocytes was discovered in leukemia and cancer patients by the precipitation method in the agar on a great amount of material.

The precipitation reaction in the agar was also success-

fully used for the analysis of the antigenic structure of the Rous sarcoma (A. I. Gusev). It was shown that there is a specific precipitating antigen in the Rous sarcoma which is absent in the normal organs of chickens and which differs from the specific antigen of the transplantable chicken sarcoma MX-659. The antigen is a tissue antigen different from the virus of the Rous sarcoma. For the virus of this sarcoma there form antibodies producing a neutralization reaction and not reacting with this tissue antigen.

Serums of cancerous animals were found to be different from the serums of healthy animals in respect to the antigens. This was shown by V. Ya. Shevlyagin when he compared the serums of rabbits having the Brown-Pierce tumors with those of healthy rabbits in the anaphylaxis reaction with desensitization in vitro. However, the specific nature of this reaction needs further study, because cross reactions between the serums of rabbits affected by the Brown-Pearce tumors and the Shope cancer have been observed. These reactions were absent in the serums of rabbits in the presence of papillomas, the Arthus phenomenon and acute inflammation processes. The anaphylaxis reaction with desensitization in vitro is very sensitive. It was successfully used by our colleague, G. T. Patrikeev (1952), for the analysis of organ specificity of animal tissues and was later suggested by Macari (1955) for the detection of a specific antigen in the serum of cancer patients.

Studies conducted in the preceding years (R. M. Radzhikhovskaya, Z. L. Baidakova, O. M. Lezhneva) have shown a possibility of the preservation of the specific antigen in tumor cells lysed with antitumor serums and the ability of these formalinized lysates to immunize against the transplanted tumors and to a certain extent against the primary tumors of animals (experiments with induced sarcomas of inbred mice, naturally developing cancer in mice of the A-strain, Brown-Pearce tumor in rabbits, Rous sarcoma, and others).

Along with the study of antitumor immunity, studies were done on the suppression of resistance in transplanting tumors dependent on the isoantigenic difference (Z. L. Baidakova). The rat sarcoma M-1 transplanted to non-inbred rats and originally induced with benzpyrene (a stock of the L. M. Shaboda Laboratory) could not be successfully transplanted to the Wistar rats, but if these rats were previously treated with the extracts of sarcoma M-1, then the



inoculations were successful and the rats died from this tumor. Such an increased susceptibility continued for only a few days (usually from the 6-7 to 14-15 day after the introduction of the extract of M-1 sarcoma), and then it disappeared. An increased susceptibility was also achieved by the introduction into the Wistar rats of the serum of rats of the same strain which had been intensively immunized with the extracts of M-1 sarcoma.

We have also studied the changes in the antigenic structure of the cells occurring under the influence of ionizing radiation (V. A. Artamonova). In the appropriate experiments, clear antigenic changes were discovered in the liver and the spleen of irradiated rabbits. Independent from us, analogous data were obtained by R. V. Petrov and L. I. Il'yina (1956). In a number of experiments, organs of the same animal (a kidney and part of the liver) were studied before and after irradiation in order to exclude a possible influence of isoantigenic differences. No changes were found in the kidney. In the liver, they were observed in most of the experiments. But sometimes changes were observed after a surgical interference without the action of irradiation. These studies should be repeated on pure-strain animals with the use of various immunological methods.

We tried to use the phenomenon of immunological rapprochement, which was studied for the first time by Hasek (1953), Medawar and their associates (1953), for obtaining serums reacting selectively with the tumor tissue (I. N. Kyukova, R. M. Radzikhovskaya, N. V. Nartsissov, T. I. Biryulina). Rats and rabbits were treated during their embryonic life with the extracts of normal tissues, and, during their adult state, they were immunized with the extracts of tumor tissues. Experiments were conducted with the tumors of animals and human beings. The serums were tested in the complement fixation reaction and in the precipitation reaction in the agar (in the last experiment).

These studies have shown that the introduction of the extracts of various tissues into the embryos of rats and rabbits with a subsequent introduction into them of the same material after birth results in a considerable suppression of the formation of antibodies during the immunization of these animals in their adult state. A suppression of the formation of antibodies was also observed if animals which were tolerant to the normal tissue were immunized with

tumor tissue. However, it was found impossible (with a few exceptions) to differentiate in this manner the extracts of the tumor and normal tissues of human beings.

Different results were obtained in analogous experiments with tumor tissues of animals. With the serums of animals which were tolerant to normal chicken tissues and which were immunized with the extracts of chicken sarcoma, it was possible to differentiate the sarcoma tissue from the tissue of a normal muscle. However, similar experiments with the cancer of mammary gland and hepatoma of mice did not produce results.

The method of artificial immunological tolerance was also used in the study of the heterotransplantation of tumors. A suspension of adenocarcinoma cells of strain mice was introduced into the embryos of the Wistar strain rats. The baby rats which grew from these embryos were inoculated hypodermically with the same tumor when they were one week and two weeks old. In this manner it was possible to obtain in rats an acclimatization of a strain-specific tumor of mice with a subsequent death of one half of the baby rats from this tumor. The tumor was passed into the baby rats three times.

One of the segments of our work is connected with the study of the pathogenic potencies of tumor-producing viruses (I. N. Kryukova, N. V. Nartissov, T. I. Biryulina). In rats infected with the Rous virus, we observed, in the embryonic stage, the development of the hemorrhagic disease with the affection of blood vessels and the formation of large cysts filled at first with a serous and then with a hemorrhagic content. We were unable to separate the virus from these affections, however, G. Ya. Svet-Moldavskiy and A. S. Skorikova succeeded in this (1958), when they made similar observations independent from us. A formalinized virus or a virus neutralized with an antiserum did not cause any pathological phenomena when it was introduced into the embryo of rats, as well as an active virus when introduced into adult rats.

In some rats which recovered after the hemorrhagic disease, tumors developed after long periods of time since the introduction of the virus (about one year). The study of these tumors is in progress. Analogous experiments were done on rabbits. The rous virus introduced into newborn baby rabbits caused a diffused fibromatosis which progres-



sed benignantly and ended with recovery. In none of these cases the virus was separated from the fibrous nodes which had formed.

All these experiments indicate that the same virus can cause different affections, both of the infectious and the neoplastic type, depending on the state and the reaction of the macro-organism. Apparently, the same factors influence to a considerable degree the separability of the virus and its masking. The phenomenon of masking was specially studied on the Shope's virus model adapted to domestic rabbits at the biological station at Sukhumi (V. A. Artamonova). This stock does not become masked when injected into domestic rabbits and can be detected in most of the papillomas, but not in a high titer (not higher than 1:100). During the malignization of the papillomas, the virus becomes completely masked. It was found that a 30 minute contact of the virus in vitro with an extract of a tumor tissue which originated from a papilloma is sufficient for the virus to lose completely its pathogenicity and not to show in the mixture. The virus processed in this manner would even lose its immunization ability, while the virus neutralized by the antibody retained it (Z. A. Postnikova). It was established (V. A. Artamonova) that when cancer tissue is fractionated, the ability to block the virus is connected with the fractions of the tumor tissue which settle at 30 and 50 percent saturation with ammonium sulfate. At present we are attempting to separate the active blocking substance from these fractions.

The etiological part of our research was dedicated to an electron-microscopic study of virus-like particles detected in tumors by various researchers and attempting to reproduce the cancer disease of human beings in animals.

As it is known, virus-like particles have been detected in many tumors of animals and human beings, whose virus etiology has not been proven. However, similar particles can also be found in normal tissues. Only in very rare cases particles of a peculiar shape were observed (for example, in human leukemia V. A. Parnes observed "caudate" particles). We attempted to obtain serological data for the characteristics of these particles (A. I. Gusev) and to study under an electronmicroscope the agglutination of these particles by specific serums. However, the agglutination titers of antisera exhausted by the particles from normal tissues were insignificant (1:40) and did not make it possible to make defi-

nite conclusions.

The attempts to transfer cancer disease of human beings to animals are successful, so far, only in respect to human leukemia. By means of introducing various leukemia material taken from leukemia patients or people who died from it into newborn mice of non-leukemia strains, it was possible to cause a transplantable leukemia in mice (V. A. Parnes and V. V. Suntsova). Analogous experiments, but with more detail and based on a greater amount of material, were set up by V. M. Bergol'tz under my direction at the Institute imeni P. A. Gertsen. In his experiments, leukemias in mice were in 34 percent of cases caused by noncellular extracts from human organs affected by leukemia. Formalin and heating at 70° in the course of one hour almost completely destroyed the activity of the factor contained in the extracts which was studied in many other respects. All these studies are in agreement with the data of Magrassi, Rzhiman and Veselyi, and other researchers who were successful in transferring human leukemia to guinea pigs and rats.

At the present time these studies have been considerably broadened (V. A. Parnes, Z. A. Postnikova, I. E. Shustrova, V. Ya. Shevlyagin, I. S. Irlin). They are conducted in the following manner: material taken from a human being is first inoculated in the cultures of a monolayer tissue, and the resistance of these cultures during a part of the experiments is weakened by X-ray irradiation. After a number of subinoculations, the cultures are introduced into the animals which were irradiated and treated with cortisone. Various methods of infecting are used, including the intratracheal method (A. M. Gardash'yan). Studies are also being planned which will take into account the possibility of the presence of a virus component having the form of nucleic acid in human tumors.

We should note the work of the development of inbred strains of mice and rats which was conducted by N. N. Medvedev. A study of the susceptibility to lung tumors of the C<sub>57</sub> and CC<sub>57</sub> strains, which were developed earlier, is now in progress.

The above studies are only the basic trends of the experimental work of our Division, which can give an idea of our goals and means of their achievement.

Active Member of the AMS USSR  
Professor L. A. Zil'ber

PRINCIPLE EXPERIMENTAL WORKS OF THE DIVISION OF  
IMMUNOLOGY AND ONCOLOGY OF THE INSTITUTE IMENI  
GAMALEYA FOR 1957-1959

- Abelev, G. I., Modification of the Precipitation Method in the Agar for Comparing 2 Antigen-Antiserum Systems. Folia biologica, 1950, No. 1 (in press).
- Abelev, G. I.; Avenirova, Z. A.; Engel'gardt, N. V.; Baidakova, Z. L., and Stepanchenok-Rudnik, G. I., Organ-specific Antigen of Liver which is Absent in the Hepatoma. Doklady AN USSR, 1959, 124, 5.
- Abelev, G. I.; Avenirova, Z. A.; Tsvetkov, V. S., Separation and Purification of the Organ-specific Antigen of Liver which is Absent in the Hepatoma. Voprosy onkologii (in press).
- Avenirova, Z. A.; Abelev, G. I., Separation of Antibodies for Specific Antigens of the Liver and Hepatoma of Mice. Problemy onkologii (in press).
- Artamonova, V. A., A Further Study of the Problem of the Influence of Ionizing Radiation on the Antigenic Properties of Proteins. Meditinskaya radiologiya, 1959 (in press).
- Artamonova, V. A.; and Levina, D. M., A Study of Antigenic Properties of Some Protein Fractions of Tumors in Strain Mice. Voprosy onkologii, 1959, 5 (1), 29.
- Baidakova, Z. L., Transplantation of Certain Tumors from Nonpedigreed Rats to Strain Rats. Neoplasma, 1959 (in press).
- Gardash'yan, A. M., Sensitivity Limit of the Anaphylaxis Reaction with Desensitization. Zhurnal mikrobiologii, epidemiologii i immunologii, 1951, 1, 71.
- Gardash'yan, A. M., and Radzikhovskaya, R. M., Detection of Antibodies in RSK in Rabbits Immunized with Various Types of Vaccines from the Brown-Pearce Tumor. Voprosy onkologii, 1958, 4 (6), 655.
- Gusev, A. I., Controls in the Electronic Microscopic Research. Voprosy virusologii, 1959, 3, 373.

- Gusev, A. I., Detection of the Specific Antigen of the Rous Sarcoma by the Precipitation Reaction in the Agar. Bulleten' eksperimental'noy biologii i meditsiny, 1959 (in press).
- Gusev, A. I., Comparative Characteristic of Specific Antigens of the Rous Sarcoma and Transplantable Methyl Cholenthrane Sarcoma of Chickens in the Precipitation Reaction in the Agar. Bull. eksperimental'noy biologii i meditsiny, 1959 (in press).
- Zil'ber, L. A., Viral Theory of the Etiology of Tumors. Patol. fiz. i eksp. terapiya, 1958, 3, 3.
- Zil'ber, L. A., A Study of the Ways of the Spreading of Cancer Diseases. Vestnik AMN SSSR, 1959, 2, 15.
- Zil'ber, L. A., Studies of Tumor Antigens. Voprosy onkologii, 1959, 5 (3), 265.
- Zil'ber, L. A., Abelev, G. I., Avenirova, Z. A., Engel'gardt, N. V., and Baidakova, Z. L., Differences in the Antigenic Structure of Cytoplasmic Granules of the Liver and the Hepatoma of Mice. Doklady AN SSSR, 1959, 124, 4.
- Zil'ber, L. A., and Artamonova, V. A., The Nature of Changes in the Antigenic Structure of Proteins under the Action of Ionizing Radiation. Meditsinskaya radiologiya, 1959, 5, 3.
- Zil'ber, L. A., Biryulina, T. I., and Nartsissov, N. V., The Reaction of Passive Hemagglutination and Its Inhibition in Chicken Sarcoma. Zhurnal gig., epid., mikrob. i immun. (Prague), 1957, 1, 68.
- Zil'ber, L. A., Gardash'yan, A. M., and Avenirova, Z. A., Peculiarities of the Antigenic Structure of Immune Globulins. Collected Articles "Sovrem. problemy immunologii", 1959, 149.
- Zil'ber, L. A., and Kryukova, I. N., Fibromatosis in Rabbits Caused by the Rous Virus, Voprosy virusologii, 1958, 3, 166.
- Zil'ber, L. A., and Kryukova, I. N., Hemorrhagic Disease in Rats Caused by the Rous Virus. Voprosy virusologii.

- gii, 1957, 4, 239.
- Zil'ber, L. A.; Nartsissov, N. V.; Biryulina, T. I., and Kryukova, I. N., Serological Differentiation of the Rous Sarcoma Extracts and the Normal Tissue. Voprosy onkologii, 1959, 4 (3), 268.
- Zil'ber, L. A., Studies of Tumor Antigens. J. Nat. Canc. Inst., 18, 1957, (3), 341.
- Zilber, L. A., Immunologike sblizeni v onkologikem vyskumu. Ceskoslovenska Biologie, 1958, VII, 4, 278.
- Zilber, L. A., Specific Tumor Antigens. Advances in Cancer Research, 1958, 5, 291.
- Zilber, L. A.; Biryulina, T. I., and Nartsissov, N. V., The Reaction of Passive Haemagglutination and Its Inhibition by Chicken Rous Sarcoma. J. Hyg. Epid. Microbiol. and Immunol., 1957, 1, 80.
- Kryukova, I. N., Acquired Tolerance to Foreign Serum in Rats. Bull. eksp. biol. i med., 1957, 4, 78.
- Kryukova, I. N., The Role of a Combined Influence of the Cell and Humoral Factors in the Destruction of Tumors. Voprosy onkologii, 1958, 4 (2), 131.
- Kryukova, I. N., A Further Study of the Hemorrhagic Disease of Rats Caused by the Rous Virus. Voprosy virusologii, 1959, 4, 486.
- Kryukova, I. N., and Lezhneva, O. M., An Attempt to Cultivate the Viruses of the Rous Sarcoma and the Shope Papilloma in the Ehrlich Ascitic Tumor and in Sarcoma No. 1. Voprosy onkologii, 1959, 5 (7), 3.
- Levina, D. M., and Artamonova, V. A., Studies on the Antigenic Properties of Certain Corpuscular and Soluble Fractions from Tumors of Inbred Mice. Bull. eksp. biol. i med., 1958, 8, 77.
- Lyudogovskaya, L. A.; Morgunova, T. D., and Postnikova, Z. A., Adsorption of the Specific Tumor Antigen and Erythrocytes. Collected Articles: "Voprosy etiologii i patogeneza opukholey", 1957, 153.



- Lyudogovskaya, L. A.; Gusev, A. I.; Abelev, G. I., The Reaction of Passive Hemagglutination in the Case of Human Tumors. Trudy 2-y Vsesoyuznoy konferentsii onkologov, 1959, 58.
- Medvedev, N. N., The CC<sub>57</sub> Strain White Mice. Doklady AN SSSR, 1958, 119 (2), 369.
- Medvedev, N. N., Oncological Characteristics of the CC<sub>57</sub> Strain Brown Mice. Bull. eksp. biol. i med., 1958, 45 (5), 104.
- Medvedev, N. N., Genetics-Dependent Differences in the Susceptibility of CC<sub>57</sub> White Mice and CC<sub>57</sub> Brown Mice to Spontaneous and Induced Lung Tumors. Doklady AN SSSR, 1958, 119 (3), 594.
- Medvedev, N. N., Blastomogenous Activity of the Virus of the Mammary Gland Cancer of Mice after 6 Years of Storing in a Dry State. Bull. eksp. biol. i med., 1958, 46 (7), 88.
- Medvedev, N. N., Genetics of Transplanted Tumors, Uspekhi sovr. biol. (in press).
- Medvedev, N. N., The Breeding of the Laboratory Strain Mice. Bull. Moskovskogo obshchestva ispyt. prirody, 1958, 63 (6), 117.
- Medvedev, N. N., CC<sub>57</sub>BR and CC<sub>57</sub>W Strains of Mice. Mouse News Letter, 1958, 18, 57.
- Medvedev, N. N., The Further Data on the Oncological Characteristics of the CC<sub>57</sub>W and CC<sub>57</sub>BR Strains of Mice. Mouse News Letter, 1958, 18, 59.
- Medvedev, N. N., 15-year Experience in Working with Strain Mice in Experimental Oncology. Patol. fiziol. i eksper. terapiya, 1959, 3 (4), 76.
- Morgunova, T. D., Postnikova, Z. A., The Mechanism of the Adsorption of Tumor-Producing Viruses by Erythrocytes. Trudy 2-y Vsesoyuznoy konferentsii onkologov, 1959, 63.
- Nartsissov, N. V.; Avenirova, Z. A.; Stepanchenok, G. J.; Solovyeva, N. J., Shope Rabbit Papilloma; Serological

and Biological Suspensions. Neoplasma, 1957, IV, 3, 196.

Nartsissov, N. V.; Biryulina, T. I.; Gusev, A. I., The Antigen of Tumors Causing Passive Hemagglutination. Trudy 2-y Vsesoyuznoy konferentsii onkologov, 1959.

Nartsissov, N. V.; Abelev, G. I., Complement Fixation Test in Rats with Primary Sarcomas Induced with Carcinogens. Neoplasma (in press).

Parnes, V. A., Analysis of the Antigens of Spleen Tissue in a Normal State and during Leukemia. Problemy gematologii i perelivaniya krovi, 1958, 6, 11.

Parnes, V. A., Immunological Study of Pathologically Changed Tissues. Voprosy onkologii, 1957, 3 (2), 179.

Parnes, V. A. and Lacour, F., Differences in the Antigenic Properties of Normal Erythrocytes and Those Affected by Certain Forms of Neoplasia. Doklady AN SSSR, 1958, 119 (2).

Postnikova, Z. A., The Methods of Intracardiac Injections for Newborn Mice and Rats. Voprosy virusologii (in press).

Radzikhovskaya, R. M., Experimental Materials on the Study of the Role of Antibodies in Immunity. Bull. eksp. biol. i med., 1958, 5, 89.

Shevlyagin, V. Ya., Detection of Antigens by Means of Fluorescent Antibodies. Uspekhi sovr. biologii, 1958, XIV (2), 218.

Shevlyagin, V. Ya., Specificity of the Antigens Found in the Blood Plasma of Rabbits with the Brown-Pearce Carcinoma. Voprosy onkologii, 1959, 5, 8, 149.

Shevlyagin, V. Ya., and Shustrova, I. E., Antigenic Peculiarities of the Plasma of Human Beings and Animals Suffering from Cancer. Trudy 2-y Vsesoyuznoy konferentsii onkologov, 1959, 82.

Shustrova, I. E., Virus-like Formations of Certain Papil-

lomas in Human Beings. Voprosy virusologii, 1959,  
2, 226.

Shershul'skaya, L. V., and Shustrova, I. E., Immunologi-  
cal Differentiation of Specific Tumor Antigens in  
Human Beings. Trudy Vsesoyuznoy konferentsii onko-  
logov, 1959, 98.

END